

Procellosaurinus erythrocerus Rodrigues, 1991 (Squamata: Gymnophthalmidae): Distribution extension

Fagner Ribeiro Delfim^{1,2*}, Daniel Oliveira Mesquita^{1,2}, Hugo Fernandes-Ferreira^{1,2}, Lucas Barbosa de Queiroga Cavalcanti¹

1 Universidade Federal da Paraíba, Centro de Ciências Exatas e da Natureza, Departamento de Sistemática e Ecologia. CEP 58051-900. João Pessoa, PB, Brazil.

2 Programa de Pós-graduação em Ciências Biológicas – Zoologia, Universidade Federal da Paraíba, Centro de Ciências Exatas e da Natureza, Departamento de Sistemática e Ecologia. CEP 58051-900. João Pessoa, PB, Brazil.

* Corresponding author. E-mail: frdelfim@yahoo.com.br

ABSTRACT: We report the northernmost occurrence of *Procellosaurinus erythrocerus* in South America. This species was originally described from the paleoquaternary sand dunes complex of the São Francisco River, and found later at Serra das Confusões, Piauí state. We found *P. erythrocerus* at São Francisco de Assis do Piauí and Pavussu municipalities, Piauí state, about 253 kilometers northwest from its type locality. This record corroborates the theory that sand dune refuges were created during recent geological eras.

The family Gymnophthalmidae is a group of Neotropical small lizards distributed between the Tehuantepec Isthmus, in southern Mexico, and northern Argentina, as well as the Caribbean and some continental islands of Central and South America. They are found in open and forested habitats, and are highly associated with the undergrowth, leaf litter and humus layer (Vanzolini *et al.* 1980; Rodrigues *et al.* 2001).

Since Gymnophthalmidae was classified as distinct from Teiidae, its systematics have been marked by the discovery of numerous genera and species (Cunha *et al.* 1991; Rodrigues 1991a, b, c; 1996a, b; 1997; Vanzolini and Carvalho 1991; Hoogmoed and Ávila-Pires 1992; Carvalho 1997; Ávila-Pires and Vitt 1998; MacCulloch and Lathrop 2001; Pellegrino *et al.* 2001; Rodrigues *et al.* 2001, 2002a, b; 2009a, b), especially with the use of cytogenetic and molecular techniques (Cole *et al.* 1993; Yonenaga-Yassuda *et al.* 1995; Pellegrino *et al.* 1999a, b; 2001; Yonenaga-Yassuda and Rodrigues 1999; Bertolloto *et al.* 2002; Castoe *et al.* 2004).

Many of these species were discovered in the Brazilian semi-arid morphoclimatic domain known as Caatinga. Within this domain, we focus on the ecoregion of the paleoquaternary dune fields of the middle São Francisco River, (Rodrigues 2003; Borges-Nojosa and Cascon 2005), the type locality of seven gymnophthalmids that were previously considered endemics (Rodrigues 1996b).

Rodrigues (1991) described two species of the genus *Procellosaurinus*. The original distribution of these two species was at two sand dune fields: *P. erythrocerus* (Figure 1) at Ibiraba (10°48' S, 42°50' W) and Alagoado, Brazilian state of Bahia (09°29' S, 41°21' W), while *P. tetradactylus* were found only at the second location. These localities are relatively isolated and harbor other endemic species of lizards and snakes (Rodrigues 1991a; 1996b). *Procellosaurinus erythrocerus* was later found at Serra das Confusões, Piauí (09°15' S, 43°32' W) (Freitas and Silva, 2007).

During a field expedition to Piauí state, we collected (SisBio/ICMBio permits n. 15363-1) *P. erythrocerus* in two municipalities: Pavussu (07°57'07" S, 43°32'01" W) and São Francisco de Assis do Piauí (08°03'37" S, 41°31'16" W) (Figure 2). The lizards were found in a forested Caatinga, with sandy soils, but no sand dunes. These records extend the distribution 253 km from the type locality and 110 km from Serra das Confusões. The specimens were deposited in the Coleção Herpetológica da Universidade Federal da Paraíba, Paraíba state, Brazil (CHUFPB 00279; CHUFPB 00932; CHUFP 00933; CHUFPB 00934; CHUFP 00968).

Other distribution extensions of gymnophthalmids from the dunes complex of the São Francisco were previously published. *Calyptommatus confusionibus* was described from Parque Nacional Serra das Confusões, Piauí state, extending the genus distribution 120 kilometers north from the Bahia sand dunes (Rodrigues *et al.* 2001). Delfim *et al.* (2006) reported *Psilophthalmus paeminus* (originally considered an endemic of the dunes), at the municipality of Canidé do São Francisco, state of Sergipe, about 540 kilometers east of the type locality.

This new report of *P. erythrocerus* supports the



FIGURE 1. *Procellosaurinus erythrocerus* collected at the dune fields of São Francisco river, state of Bahia, Brazil.

hypothesis suggested by Rodrigues *et al.* (2001), which posits that these gymnophthalmid distributions are based on the recent geological history (Pleistocene-Holocene) of the São Francisco sand dunes. Southeast winds may have affected the distribution and size of dune fields (Ab'Saber 1977; Rodrigues 1987; Oliveira *et al.* 1999), and consequently the distribution of *P. erythrocerus*. The most recent retraction of the dunes restricted them to their present distribution and size, creating some sandy "islands" which acted as refuges to species associated with them. This interpretation is similar to the Pleistocene forest refuges suggested by Vanzolini and Williams (1981).

Additionally, the increasing knowledge of the Caatinga domain has led to the distribution extension of some species previously described as endemic to the São Francisco River dunes. These species are restricted to sandy areas, so all of those areas could have been an extensive sand dune field in the past.

Finally, the presence of *P. erythrocerus* in two localities of Piauí state presents a significant distribution extension from previous known localities, and the possible presence of another species, *P. tetradactylus*, should be investigated in Piauí.




FIGURE 2. Known geographic distribution of *Procellosaurinus erythrocerus*, including the new localities reported here. Circles = previous localities: 1) Alagoado; 2) Ibiraba; 3) Serra das Confusões; triangles = new records: 4) São Francisco de Assis do Piauí; 5) Pavussu.

ACKNOWLEDGMENTS: We are grateful to Bruno Campos, Gustavo Toledo and Luciana Lobo, who provided valuable assistance with data collection. We thank Stephen Ferrari for his revision of the text and Anderson Feijó for the production of the map. We also thank Miguel T. Rodrigues for the

P. erythrocerus photo. This work is supported by research fellowship from Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq to Daniel Oliveira Mesquita, a doctorate fellowship from CNPq to Fagner Ribeiro Delfim, a scientific initiation to Lucas Barbosa de Queiroga Cavalcanti, and a doctorate fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - CAPES to Hugo Fernandes-Ferreira.

LITERATURE CITED

- Ab'Saber, A.N. 1977. Espaços ocupados pela expansão dos climas secos na América do Sul, por ocasião dos períodos glaciais quaternários. *Paleoclimas* 3: 1-19.
- Ávila-Pires, T.C.S. and L.J. Vitt. 1998. A new species of *Neusticurus* (Reptilia: Gymnophthalmidae) from the Rio Juruá, Acre, Brazil. *Herpetologica* 54(2): 235-245.
- Bertolotto, C.E.V., K.C.M. Pellegrino, M.T. Rodrigues and Y. Yonenaga-Yassuda. 2002. Comparative cytogenetics and supernumerary chromosomes in the Brazilian lizard genus *Enyalus* (Squamata, Polychrotidae). *Hereditas* 136: 51-57.
- Borges-Nojosa, D.M. and P. Cascon. 2005. Herpetofauna da área Reserva da Serra das Almas, Ceará; p. 245-260. In F.S. Araújo, M.J.N. Rodal and M.R.V. Barbosa (org.). *Análise das variações da biodiversidade do Bioma Caatinga: suporte a estratégias regionais de conservação*. Brasília: Ministério do Meio Ambiente.
- Carvalho, C.M. 1997. Uma nova espécie de microteídeo do gênero *Gymnophthalmus* do Estado de Roraima, Brasil (Sauria, Gymnophthalmidae). *Papéis Avulsos de Zoologia* 40(10): 163-176.
- Castoe, T.A., T.M. Doan and C.L. Parkinson. 2004. Data partitions and complex models in Bayesian analysis: the phylogeny of Gymnophthalmid lizards. *Systematic Biology* 53(3): 448-469.
- Cole, C.J., H.C. Dessauer and A.L. Markezich. 1993. Missing Link Found: The Second Ancestor of *Gymnophthalmus underwoodi* (Squamata: Teiidae), A South American Unisexual Lizard of Hybrid Origin. *American Museum Novitates* 3055: 1-13.
- Cunha, O.R., J.S. Lima-Verde and A.C. Lima. 1991. Novo gênero e espécie de lagarto do Estado do Ceará (Lacertilia: Teiidae). *Boletim do Museu Paraense Emílio Goeldi* 7(2): 163-176.
- Delfim, F.R., E.M. Gonçalves and S. T. Silva. 2006. Squamata, Gymnophthalmidae, *Psilophthalmus paeminus*: Distribution extension, new state record. *Check List* 2(3): 89-92.
- Freitas, M.A. and T.F.S. Silva. 2007. *A Herpetofauna das Caatingas e Áreas de Altitude do Nordeste Brasileiro*. Pelotas: União Sulamericana de Estudos da Biodiversidade. 388 p.
- Hoogmoed, M.S. and T.C.S. Ávila-Pires. 1992. Studies on the species of the South American lizard genus *Arthrosaura* Boulenger (Reptilia: Sauria: Teiidae), with the resurrection of two species. *Zoologische Mededelingen* 66: 453-484.
- MacCulloch, R.D. and A. Lathrop. 2001. A new species of *Arthrosaura* (Sauria: Teiidae) from the highlands of Guyana. *Caribbean Journal of Science* 37(3-4): 174-181.
- Oliveira, P.E., A.M.F. Barreto and K. Suguio. 1999. Late Pleistocene/Holocene climatic and vegetational history of the Brazilian Caatinga: the fossil dunes of the middle São Francisco River. *Palaeogeography, palaeoclimatology, palaeoecology* 152: 319-337.
- Pellegrino, K.C.M., M.T. Rodrigues and Y. Yonenaga-Yassuda. 1999a. Chromosomal polymorphisms due to supernumerary chromosomes and pericentric inversions in the eyelidless microteiid lizard *Notobachia ablephara* (Squamata, Gymnophthalmidae). *Chromosome Research* 7: 247-259.
- Pellegrino, K.C.M., M.T. Rodrigues and Y. Yonenaga-Yassuda. 1999b. Chromosomal evolution in the Brazilian lizards of genus *Leposoma* (Squamata, Gymnophthalmidae) from Amazon and Atlantic rain forests: banding patterns and FISH of telomeric sequences. *Hereditas* 131: 5-21.
- Pellegrino, K.C.M., M.T. Rodrigues, Y. Yonenaga-Yassuda and J.W. Sites, Jr. 2001. A molecular perspective on the evolution of microteiid lizards (Squamata, Gymnophthalmidae), and a new classification for the family. *Biological Journal of the Linnean Society* 74: 315-338.
- Rodrigues, M.T. 1987. Sistemática, Ecologia e Zoogeografia dos *Tropidurus* do Grupo *Torquatus* ao Sul do Rio Amazonas (Sauria, Iguanidae). *Arquivos de Zoologia* 31(3): 105 - 231.
- Rodrigues, M.T. 1991a. Herpetofauna das dunas interiores do Rio São Francisco: Bahia: Brasil: I. Introdução à área e descrição de um novo gênero de microteídeos (*Calyptommatius*) com notas sobre sua ecologia, distribuição e especiação (Sauria, Teiidae). *Papéis Avulsos de Zoologia* 37(19): 285-320.
- Rodrigues, M.T. 1991b. Herpetofauna das dunas interiores do Rio São Francisco: Bahia: Brasil: II. *Psilophthalmus*: um novo gênero de microteídeos sem pálpebras (Sauria, Teiidae). *Papéis Avulsos de Zoologia* 37(20): 321-327.

- Rodrigues, M.T. 1991c. Herpetofauna das dunas interiores do Rio São Francisco: Bahia: Brasil: III. *Procellosaurinus*: um novo gênero de microteídeos sem pálpebras, com a redefinição do gênero *Gymnophthalmus* (Sauria, Teiidae). *Papéis Avulsos de Zoologia* 37(21): 329-342.
- Rodrigues, M.T. 1996a. A new species of lizard, genus *Micrablepharus* (Squamata: Gymnophthalmidae), from Brazil. *Herpetologica* 52(4): 535-541.
- Rodrigues, M.T. 1996b. Lizards, snakes, and amphisbaenians from the quaternary sand dunes of the middle Rio São Francisco, Bahia, Brazil. *Journal of Herpetology* 30(4): 513-523.
- Rodrigues, M.T. 1997. A new species of *Leposoma* (Squamata, Gymnophthalmidae) from the Atlantic Forest of Brazil. *Herpetologica* 53(3): 383-389.
- Rodrigues, M.T. 2003. Herpetofauna da Caatinga; p. 181-236. In I.R. Leal, M. Tabarelli and J.M.C. Silva (ed.). *Ecologia e Conservação da Caatinga*. Recife: Ed. Universitária-UFPE.
- Rodrigues, M.T., H. Zaher and F. Curcio. 2001. A new species of lizard, genus *Calyptommatus*, from the Caatingas of the state of Piauí, Northeastern Brazil (Squamata, Gymnophthalmidae). *Papéis Avulsos de Zoologia* 41(28): 529-546.
- Rodrigues, M.T., M. Dixo, D. Pavan and V.K. Verdade. 2002a. A new species of *Leposoma* (Squamata, Gymnophthalmidae) from the remnant Atlantic Forests of the State of Bahia, Brazil. *Papéis Avulsos de Zoologia* 42(14): 335-350.
- Rodrigues, M.T., M. Dixo and G.M. Accacio. 2002b. A large sample of *Leposoma* (Squamata, Gymnophthalmidae) from the Atlantic Forests of Bahia, the status of *Leposoma annectans* Ruibal, 1952, and notes on conservation. *Papéis Avulsos de Zoologia* 42(5): 103-117.
- Rodrigues, M.T., J. Cassimiro, M.A. Freitas and T.F.S. Silva. 2009a. A new microteiid lizard of the genus *Acratosaura* (Squamata: Gymnophthalmidae) from Serra do Sincorá, State of Bahia, Brazil. *Zootaxa* 2013: 17-29.
- Rodrigues, M.T., M.A. Freitas and T.F.S. Silva. 2009b. New species of earless lizard genus *Heterodactylus* (Squamata: Gymnophthalmidae) from the highlands of Chapada Diamantina, State of Bahia, Brazil. *Journal of Herpetology* 43: 605-611.
- Vanzolini, P.E. and C.M. Carvalho. 1991. Two sibling and sympatric species of *Gymnophthalmus* in Roraima, Brasil (Sauria, Teiidae). *Papéis Avulsos de Zoologia* 37(12): 173-226.
- Vanzolini, P. E. and E.E. Williams. 1981. The Vanishing Refuge: a mechanism for ecogeographic speciation. *Papéis Avulsos de Zoologia* 34(23): 251-255.
- Vanzolini, P.E., A.M.M. Ramos-Costa and L.J. Vitt. 1980. *Répteis das Caatingas*. Rio de Janeiro: Academia Brasileira de Ciências. 161 p.
- Yonenaga-Yassuda, Y. and M.T. Rodrigues. 1999. Supernumerary chromosome variation, heteromorphic sex chromosomes and banding patterns in microteiid lizards of the genus *Micrablepharus* (Squamata, Gymnophthalmidae). *Chromosome Research* 7: 247: 259.
- Yonenaga-Yassuda, Y., P.E. Vanzolini, M.T. Rodrigues and C.M. Carvalho. 1995. Chromosome banding patterns in the unisexual microteiid *Gymnophthalmus underwoodi* and in two related sibling species (Gymnophthalmidae, Sauria). *Cytogenetic and Cell Genetics* 70: 29-34.

RECEIVED: October 2011

LAST REVISED: November 2011

ACCEPTED: November 2011

PUBLISHED ONLINE: December 2011

EDITORIAL RESPONSIBILITY: Ross MacCulloch